

AD-A109 915

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD RALSTON (ALBERTA) F/8 6/5
PARTIAL CHARACTERIZATION OF SMALL PLASMID DEOXYRIBONUCLEIC ACID--ETC(U)
MAY 81 A R BHATTI, K O'HARA, L A WHITE


UNCLASSIFIED

NL

1-1
2-1
3-1



END
DATE
FILMED
3-82
DTIC



UNCLASSIFIED

DRES

SUFFIELD REPORT

NO. 295

AD A109915

LEVEL II

③

UNLIMITED
DISTRIBUTION
ILLIMITÉE

PARTIAL CHARACTERIZATION OF SMALL PLASMID DEOXYRIBONUCLEIC
ACID PRESENT IN *NEISSERIA MENINGITIDIS* (U)

by

A.R. Bhatti, K. O'hara, L.A. White and L.E. Bryan

PCN 16A10

DTIC
SELECTED
JAN 21 1982
E

May 1981



DEFENCE RESEARCH ESTABLISHMENT SUFFIELD : RALSTON : ALBERTA

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD
RALSTON ALBERTA

DRES ~~SUFFIELD REPORT NO~~ 295

PARTIAL CHARACTERIZATION OF SMALL PLASMID DEOXYRIBONUCLEIC
ACID PRESENT IN NEISSERIA MENINGITIDIS (4)

by

A.R. Bhatti, K. O'hara, L.A. White and L.E. Bryan

PCN 16A10

| | |
|--------------------|-------------------------------------|
| Accession For | |
| INTL. C. I. | <input checked="" type="checkbox"/> |
| PLAS. C. I. | <input type="checkbox"/> |
| UNIT. C. I. | <input type="checkbox"/> |
| J. I. | <input type="checkbox"/> |
| By | |
| Dist. | |
| Availability Codes | |
| Dist | Avail. and/or Special |
| A | |

WARNING
The use of this information is permitted subject to recognition
of proprietary and patent rights.

DTIC
COPY
INSPECTED
3

UNCLASSIFIED

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD
RALSTON ALBERTA

SUFFIELD REPORT NO. 295

PARTIAL CHARACTERIZATION OF SMALL PLASMID DEOXYRIBONUCLEIC
ACID PRESENT IN *NEISSERIA MENINGITIDIS* (U)

by

A.R. Bhatti, K. O'hara, L.A. White and L.E. Bryan

ABSTRACT

The presence of plasmid DNA in *Neisseria meningitidis* was investigated. *N. meningitidis* isolates from clinical meningitis and highly virulent in mice, were found to harbour a small plasmid. Agarose gel electrophoresis of crude lysates from different strains revealed a single plasmid DNA species with the same apparent electrophoretic mobility. Using agarose gel electrophoresis its molecular weight was calculated as 3.4 megadaltons. Electron microscopic observation revealed that plasmid DNA molecules were predominantly in the covalently closed form. The estimated contour length of the open circular form was 1.6 μm . The function of the plasmid DNA is presently unknown.

(U)

UNCLASSIFIED

UNCLASSIFIED

MILITARY SIGNIFICANCE

The discovery that virulent strains of *N. meningitidis* harbour a plasmid(s) is of great significance to military preventive medicine and biological defence. The possible role of this plasmid in virulence may account for our earlier observations at the Canadian Forces Recruit Schools of lack of correlation between carrier rates and cases of overt meningococcal disease (16). In an epidemiological sense, the fact that important virulence mediating factor(s) may be plasmid-coded could account for the patterns of recent epidemics of meningococcal meningitis in Brazil and Finland. In those outbreaks, there was a rapid increase in overt disease followed by an extended peak and equally abrupt termination of this disease, although *N. meningitidis* of apparently identical antigenic pattern was still present in the community. One can speculate that the plasmid was suddenly lost or some changed environmental factor favoured the maintenance of non-plasmid containing bacteria. Our studies, as yet unpublished, indicate that these plasmids are easily lost on laboratory subculture. Virulence mediating plasmids have also been reported in other bacterial species (4, 17-20), and have been successfully transferred to other strains of the same species (3, 4, 21) or different bacterial species (3, 22, 23) as well. It is apparent, therefore, that increased attention must be paid to the possibility that the development of a new bacterial strain of increased virulence is a distinct possibility.

Comparative studies of plasmid-harboursing and plasmid-less *N. meningitidis* strains will help in elucidating the nature of the factor(s) responsible for the virulence of pathogenic strains of this organism. Such studies will further help in the diagnosis, treatment and control of meningococcal meningitis.

UNCLASSIFIED

ACKNOWLEDGMENTS

We acknowledge the capable technical assistance of Mr. M. Knodel.

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD
RALSTON ALBERTA

SUFFIELD REPORT NO. 295

PARTIAL CHARACTERIZATION OF SMALL PLASMID DEOXYRIBONUCLEIC
ACID PRESENT IN *NEISSERIA MENINGITIDIS* (U)

by

A.R. Bhatti, K. O'hara, L.A. White and L.E. Bryan

INTRODUCTION

The presence of extra-chromosomal genetic elements (plasmids) is well documented for the members of Enterobacteriaceae, *Staphylococcus aureus* and several species of *Pseudomonas* (1-3). In the genus *Neisseria*, different kinds of plasmids have been reported and characterized from *Neisseria gonorrhoeae* (4), but little is known about the existence of plasmid DNA in *N. meningitidis*.

N. meningitidis continues to be an important etiological agent of bacteremia and meningitis in both the civilian and military population. The organism can be carried in the nasopharynx of apparently healthy individuals and this carrier state may serve as a reservoir of potentially infectious organisms. Lack of immunity usually facilitates the subsequent development of bacteremia which can lead to an acute purulent meningitis. *N. meningitidis* produces several factors, an anti-phagocytic

UNCLASSIFIED

capsule, lipopolysaccharide and pili, which have been shown to play a role in virulence in the closely-related bacterium, *N. gonorrhoeae* (5, 6). In addition, this organism has been shown to produce different protease activities (7, 8). We have recently reported the presence of plasmid DNA in a virulent strain of *N. meningitidis* (submitted for publication). The use of gel electrophoresis techniques (10) has simplified the detection of plasmids. In this communication, we describe the detection and partial characterization of plasmid DNA present in some virulent strains of *N. meningitidis*.

MATERIALS AND METHODS

Bacterial Strains.

N. meningitidis, strains M1011, DRES 30 and LCDC 604 are clinical disease isolates and DRES 03 is a carrier isolate. *Escherichia coli* K12, SA1306 pr° , met^- , nal^r (RP4), *Pseudomonas aeruginosa* 130 and *E. coli* strain F-1161 (pAF-6) (kindly supplied by Dr. T. Tamoki, University of Calgary) were used as sources of DNA standards to determine the molecular weight of the plasmid DNA of strain M1011. *N. meningitidis*, strain M1011, was cultivated in Brain Heart Infusion broth (BHI broth) supplemented with 1% (v/v) IsoVitaleX (Baltimore Biological Laboratories, Cockeysville, Md., U.S.A.) for 20 hrs at 35°C in an atmosphere of 5% CO₂ in air (9). Other bacterial strains were grown in BHI broth at 37°C for the same time.

Preparation of Cell Lysates and Agarose Gel Electrophoresis.

Crude cell lysates for plasmid DNA analysis were prepared according to published procedures (10, 11). Agarose gel electrophoresis and detection of plasmid DNA was carried out as described by Hansen and Olsen (10).

Purification of Plasmid DNA.

Purified preparations of plasmid DNA were obtained by CsCl - ethidium bromide - buoyant density centrifugation as described by Grinsted *et al.* (12). Further purification of plasmid DNA was by agarose gel

electrophoresis as described above. Fractions containing plasmid DNA were pooled and dialyzed against the TES buffer (Tris 30 mM, EDTA 5 mM, NaCl 50 mM, pH 8.0) of Meyers *et al.* (11).

Electron Microscopy of Plasmid DNA.

A modification of the aqueous technique described by Davis *et al.* (13) was used for the preparation of plasmid DNA for electron microscope examination as follows:

A grid with 0.3% formvar was prepared, coated with plasmid DNA, dipped into the cytochrome C solution for 1 to 15 min and stained with 4 times concentrated uranyl acetate solution. The plasmid DNA was examined with a Hitachi S-450 electron microscope operating at an acceleration voltage of 60 kV.

RESULTS AND DISCUSSION

N. meningitidis strain M1011, a clinical isolate has been previously shown to be highly virulent in a mouse model of meningococcal septicemia (Holbein, B.E., Infect. Immun., in press). As some virulence factors are specified by plasmid DNA (1 - 4), an intensive search for such DNA was carried out in *N. meningitidis* strain M1011. Preliminary investigations demonstrated that plasmid DNA was difficult to detect by the commonly used methods of Myers *et al.* (11). It was more readily detected when the plasmid isolation method of Hansen and Olsen (10) was used. Figure 1 shows the results of an agarose gel electrophoresis of plasmid DNA from *N. meningitidis* strain M1011 extracted by the Hansen and Olsen method (10). The plasmid band marked pB0-1 was consistently observed (Fig. 1A). Presence of plasmid DNA was further investigated in other strains of *N. meningitidis*. The results presented in Figure 2 clearly illustrate the presence of plasmid DNA in the virulent strains LCDC 604, M1011 and DRES 30 (Fig. 2; column 1 - 3); whereas, DRES 03 (Fig. 2; column 6), an avirulent strain, did not contain plasmid. These results suggest the presence of plasmid DNA may be a property of virulent strains. Using agarose gel electrophoresis (10) and plasmid DNA standards (Fig. 3) the molecular weight of pB0-1 was calculated to be 3.4

megadaltons. The plasmid DNA from strain M1011 was purified by CsCl - ethidium bromide density centrifugation as described (12). The purified plasmid pB0-1 after agarose gel electrophoresis gave a single DNA species (data not shown) corresponding to that shown in Fig. 1 for the crude cell lysate. Electron microscopic observation revealed that under standard conditions of preparation (13), plasmid pB0-1 was observed in the covalently closed circular form (Fig. 4). However, when the period of cytochrome C treatment was extended to more than 15 min, the open circular plasmid DNA form was obtained. The contour length of plasmid pB0-1 molecules was determined by electron microscopy, and plasmid DNA length with an average value of $1.6 \mu\text{m}$ was calculated. Using the standard conversion factor of $2.07 \pm 0.04 \times 10^6/\mu\text{m}$, the molecular weight was estimated to be 3.3 megadaltons. This value agrees closely with that obtained by agarose gel electrophoresis.

Although plasmid DNA has been detected in *N. gonorrhoea* (4), it has not yet been reported in strains of *N. meningitidis*. Results reported in this study demonstrate the presence of at least one type of plasmid DNA. In some experiments, in addition to pB0-1, two slower moving plasmid DNA bands (not clearly visible in the photograph) were observed after agarose gel electrophoresis (Fig. 1B). These additional bands of plasmid could be larger molecular weight plasmids or conversely might be the open circular and linear form of the 3.4 megadalton plasmid DNA.

The relationship of this plasmid DNA to virulence in *N. meningitidis* strain M1011 remains unknown. Studies are currently in progress comparing nonisogenic virulent and avirulent isolates for the presence of plasmid DNA. As well, we are currently attempting to reduce the virulence of strain M1011 by various techniques. If successful, nonvirulent derivatives and virulent isolates of *N. meningitidis* strain M1011 will be compared for plasmid DNA. At the present time, the function of pB0-1 is cryptic. Cryptic plasmids have been detected in a wide variety of bacteria. Small molecular weight plasmids detected in *N. gonorrhoea* (4) have been shown to code for β -lactamase activity. However, this is not a function of the plasmid in *N. meningitidis*.

REFERENCES

1. Novick, R.P. (1969). Extrachromosomal Inheritance in Bacteria. *Bacteriol. Rev.* 32, 210-225.
2. Clowes, R.C. (1972). Molecular Structure of Bacterial Plasmids. *Bacteriol. Rev.* 36, 361-404.
3. Chakarabarty, A.M. (1976). Plasmids in *Pseudomonas*. *Annu. Rev. Genet.* 10, 7-30.
4. Elwell, L.P., and Falkow, S. (1977). Plasmids of the Genus *Neisseria*. In *The Gonococcus*, pp. 137-154, John Wiley, New York, USA.
5. Hendley, J.W., Powell, K.R., Jordan, J.R., Rodewald, R.D., and Volk, W.A. (1978). Capsules of *Neisseria gonorrhoeae*. *Immunology of Neisseria gonorrhoeae*, pp. 116-120, Amer. Soc. Microbiol.
6. McGee, Z.A., Melly, M.A., Gregg, C.R., Horn, R.G., Taylor-Robinson, D., Johnson, A.P., and McCutchan, J.A. (1978). Virulence Factors of Gonococci: Studies Using Human Fallopian Tube Organ Cultures. *Immunology of Neisseria gonorrhoeae*, pp. 258-262, Amer. Soc. Microbiol.
7. Kilian, M., Mestecky, J., Kulhavy, R., Tomana, M., and Butler, W.T. (1980). IgA1 Proteases from *Haemophilus influenzae*, *Streptococcus pneumoniae*, *Neisseria meningitidis* and *Streptococcus sanguis*: Comparative Immunochemical. *J. Immunol.* 124, 2596-2600.
8. Chen, K.C.S., and Buchanan, T.M. (1980). Hydrolases from *Neisseria meningitidis*. The Study of Gonococsin, an Aminopeptidase-P, a Proline Aminopeptidase and Asparaginase. *J. Biol. Chem.* 225, 1704-1710.
9. Holbein, B.E., Jericho, K.W.F., and Likes, G.C. (1979). *Neisseria meningitidis* Infection in Mice: Influence of Iron, Variation in Virulence Among Strains and Pathology. *Infect. Immun.* 24, 545-551.
10. Hansen, J.B., and Olsen, R.H. (1978). Isolation of Large Bacterial Plasmids and Characterization of the P-2 to Incompatibility Group Plasmids pMG-1 and pMG-5. *J. Bacteriol.* 135, 227-238.
11. Meyers, J.A., Sanchez, D., Elwell, L.P., and Falkow, S. (1976). Simple Agarose Gel Electrophoretic Method. The Identification and Characterization of Plasmid DNA. *J. Bacteriol.* 127, 1529-1537.

REFERENCES (Cont'd)

12. Grinsted, J., Saunders, J.R., Ingram, L.C., Sykes, R.B., and Richmond, M.H. (1972). Properties of an R-Factor Which Originated in *Pseudomonas aeruginosa*. J. Bacteriol. 110, 529-537.
13. Davis, R., Simon, M., and Davidson, D. (1971). Nucleic Acids. Methods Enzymol. 21, 413-428.
14. Shahrabadi, M.S., Bryan, L.E., and Van Den Elzen, H.M. (1975). Further Properties P-2 R Factors of *Pseudomonas aeruginosa*. Can. J. Microbiol. 21, 592-605.
15. Law, S., Tamoki, T., Kruezaier, F., and Dugaiczky, A. (1980). Molecular Cloning of DNA Complementary to a Mouse Alpha Fetoprotein in RNA Sequence. Gene 10, 53-61.
16. White, L.A., Spence, M.R., and Appleford, L. (1979). Meningococcal Carriage in Canadian Forces Recruits. Proc. Can. Soc. Microbiol. 19, pp. 49.
17. Zink, D.L., Feeley, J.C., Wells, J.G., Vanderzant, C., Vickery, J.C., Roof, W.D., and O'Donovan, G.A. (1980). Plasmid-Mediated Tissue Invasiveness in *Yersinia enterocolitica*. Nature 283, 224-226.
18. Jamieson, A.F., Bremner, D.A., Bergquist, P.L., and Lane, H.E. (1979). Characterization of Plasmids Antibiotic-Resistant *Shigella* Isolates by Agarose Gel Electrophoresis. J. Gen. Microbiol. 113, 73-81.
19. Stuy, J.H. (1979). Plasmid Transfer in *Haemophilus influenzae*. J. Bacteriol. 139, 520-529.
20. Smith, H.W., and Huggins, M.B. (1976). Further Observations on the Association of the Colicine V Plasmid of *Escherichia coli* with Pathogenicity and with Survival in the Alimentary Tract. J. Gen. Microbiol. 92, 335-350.
21. Norlander, L., Davies, J., and Normark, S. (1979). Genetic Exchange Mechanisms in *Neisseria gonorrhoeae*. J. Bacteriol. 138, 756-761.
22. Smith, H.W. (1974). A Search for Transmissible Pathogenic Characters in Invasive Strains of *Escherichia coli*: The Discovery of a Plasmid-Controlled Toxin and a Plasmid-Controlled Lethal Character Closely Associated, or Identical with Colicine V. J. Gen. Microbiol. 83, 95-111.

UNCLASSIFIED

/7

REFERENCES (Cont'd)

23. Kopecko, D.J., Washington, O., and Formal, S.B. (1980). Genetic and Physical Evidence for Plasmid Control of *Shigella sonnei* from 1 Cell Surface Antigen. Infect. Immun. 29, 207-214.

UNCLASSIFIED




Figure 1. Agarose gel electrophoresis of plasmid DNA from *N. meningitidis* strain M1011.

- (A) Ethanol precipitated DNA (10 μ L) from cleared lysate of *N. meningitidis* strain M1011.
- (B) 20 μ L of sample A.
- (C) Ethanol precipitated DNA (5 μ L) from cleared lysate of *E. coli* 1306 (RP4).

UNCLASSIFIED

SR No. 295




Figure 2. An agarose gel electrophoresis of crude lysate obtained from different strains of *N. meningitidis* and *E. coli*. Columns 1, 2 and 3 represent the presence of plasmid DNA in *N. meningitidis* strains LCDC 604A, M1011, DRES 30 and absence of plasmid DNA in DRES 03, respectively. Whereas, column 4 and 5 gave the results for plasmid DNA present in *E. coli* strains SA 1306 and F-1161 used as for marker plasmid DNA.

UNCLASSIFIED




Figure 3. Molecular weight of plasmid DNA from *N. meningitidis* strain M1011. Plasmid RP4 (mol. wt. 36 megadaltons) of *E. coli* (11), R-130a (mol. wt. 23 megadaltons) of *P. aeruginosa* (14) and pAF-6 (mol. wt. 4 megadaltons) of *E. coli* F-1161 (15) were used as standard plasmids DNA for the determination of the mol. wt. of plasmid pB0-1 from *N. meningitidis* strain M1011. Symbols represent o, RP4; Δ, R-130A; □, pAF-6 and •, pB0-1, respectively.

UNCLASSIFIED

SR No. 295



Figure 4. Electron micrograph of covalently closed circular plasmid DNA molecule of *N. meningitidis* M1011. Magnification $\times 60,400$.

UNCLASSIFIED

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall document is classified)

| | | | |
|--|--|--|---------------------------|
| 1. ORIGINATING ACTIVITY DEFENCE RESEARCH ESTABLISHMENT SUFFIELD | | 2a. DOCUMENT SECURITY CLASSIFICATION UNCLASSIFIED | |
| | | 2b. GROUP | |
| 3. DOCUMENT TITLE PARTIAL CHARACTERIZATION OF SMALL PLASMID DEOXYRIBONUCLEIC ACID PRESENT IN <i>NEISSERIA MENINGITIDIS</i> (U) | | | |
| 4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Suffield Report | | | |
| 5. AUTHOR(S) (Last name, first name, middle initial) Bhatti, A.R., O'hara, K., White, L.A. and Bryan, L.E. | | | |
| 6. DOCUMENT DATE MAY 1981 | | 7a. TOTAL NO. OF PAGES 13 | 7b. NO. OF REFS 23 |
| 8a. PROJECT OR GRANT NO. PCN 16A10 | | 9a. ORIGINATOR'S DOCUMENT NUMBER(S) SUFFIELD REPORT NO. 295 | |
| 8b. CONTRACT NO. | | 9b. OTHER DOCUMENT NO.(S) (Any other numbers that may be assigned this document) | |
| 10. DISTRIBUTION STATEMENT Qualified requesters may obtain copies of this document from their defence documentation centre | | | |
| 11. SUPPLEMENTARY NOTES 11 | | 12. SPONSORING ACTIVITY | |
| 13. ABSTRACT The presence of plasmid DNA in <i>Neisseria meningitidis</i> was investigated. <i>N. meningitidis</i> isolates from clinical meningitis and highly virulent in mice, were found to harbour a small plasmid. Agarose gel electrophoresis of crude lysates from different strains revealed a single plasmid DNA species with the same apparent electrophoretic mobility. Using agarose gel electrophoresis its molecular weight was calculated as 3.4 megadaltons. Electron microscopic observation revealed that plasmid DNA molecules were predominantly in the covalently closed form. The estimated contour length of the open circular form was 1.6 μ m. The function of the plasmid DNA is presently unknown. (U) | | | |

KEY WORDS

Neisseria meningitidis
Plasmid DNA
Electron Microscope
Virulence, Electrophoresis

INSTRUCTIONS

1. **ORIGINATING ACTIVITY** Enter the name and address of the organization issuing the document.
- 2a. **DOCUMENT SECURITY CLASSIFICATION:** Enter the overall security classification of the document including special warning terms whenever applicable.
- 2b. **GROUP** Enter security reclassification group number. The three groups are defined in Appendix 'M' of the DRB Security Regulations.
3. **DOCUMENT TITLE** Enter the complete document title in all capital letters. Titles in all cases should be unclassified. If a sufficiently descriptive title cannot be selected without classification, show title classification with the usual one-capital-letter abbreviation in parentheses immediately following the title.
4. **DESCRIPTIVE NOTES** Enter the category of document, e.g. technical report, technical note or technical letter. If appropriate, enter the type of document, e.g. interim, progress, summary, annual or final. Give the inclusive dates when a specific reporting period is covered.
5. **AUTHOR(S)** Enter the name(s) of author(s) as shown on or in the document. Enter last name, first name, middle initial. If military, show rank. The name of the principal author is an absolute minimum requirement.
6. **DOCUMENT DATE** Enter the date (month, year) of Establishment approval for publication of the document.
- 7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. **NUMBER OF REFERENCES** Enter the total number of references cited in the document.
- 8a. **PROJECT OR GRANT NUMBER** If appropriate, enter the applicable research and development project or grant number under which the document was written.
- 8b. **CONTRACT NUMBER** If appropriate, enter the applicable number under which the document was written.
- 9a. **ORIGINATOR'S DOCUMENT NUMBER(S)** Enter the official document number by which the document will be identified and controlled by the originating activity. This number must be unique to this document.
- 9b. **OTHER DOCUMENT NUMBER(S)** If the document has been assigned any other document numbers (either by the originator or by the sponsor), also enter this number(s).
10. **DISTRIBUTION STATEMENT:** Enter any limitations on further dissemination of the document, other than those imposed by security classification, using standard statements such as:
 - (1) "Qualified requesters may obtain copies of this document from their defence documentation center."
 - (2) "Announcement and dissemination of this document is not authorized without prior approval from originating activity."
11. **SUPPLEMENTARY NOTES.** Use for additional explanatory notes.
12. **SPONSORING ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring the research and development. Include address.
13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document, even though it may also appear elsewhere in the body of the document itself. It is highly desirable that the abstract of classified documents be unclassified. Each paragraph of the abstract shall end with an indication of the security classification of the information in the paragraph (unless the document itself is unclassified) represented as (TS), (S), (C), (R), or (U).

The length of the abstract should be limited to 20 single-spaced standard typewritten lines: 7 1/4 inches long.
14. **KEY WORDS.** Key words are technically meaningful terms or short phrases that characterize a document and could be helpful in cataloging the document. Key words should be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context.

FILMED

3-82